





All cells are surrounded by a thin, flexible barrier known as the **cell membrane**.

Many cells also produce a strong supporting layer around the membrane known as a **cell wall**.

Copyright Pearson Prentice Hal













7-3 Cell Boundaries 📫 Cell Membrane

The lipid bilayer gives cell membranes a flexible structure that forms a barrier between the cell and its surroundings.









Cell Walls

Cell walls are found in plants, algae, fungi, and many prokaryotes.

The cell wall lies outside the cell membrane.

Most cell walls are porous enough to allow water, oxygen, carbon dioxide, and certain other substances to pass through easily.

7-3 Cell Boundaries 🔿 Diffusion Through Cell Boundaries

Diffusion Through Cell Boundaries

Every living cell exists in a liquid environment.

The cell membrane regulates movement of dissolved molecules from the liquid on one side of the membrane to the liquid on the other side.





7-3 Cell Boundaries 🛶 Diffusion Through Cell Boundaries

Diffusion

Particles in a solution tend to move from an area where they are more concentrated to an area where they are less concentrated.

This process is called **diffusion**.

When the concentration of the solute is the same throughout a system, the system has reached **equilibrium**.









Solute particles move from the side of the membrane with a higher concentration of solute to the side of the membrane with a lower concentration of solute. The solute particles will continue to diffuse across the membrane until equilibrium is

6











7-3 Cell Boundaries 📫 Osmosis

Water tends to diffuse from a highly concentrated region to a less concentrated region.

If you compare two solutions, the more concentrated solution is **hypertonic** ("above strength").

The more dilute solution is **hypotonic** ("below strength").

7-3 Cell Boundaries 📫 Osmosis

When concentrations of solutions are the same on both sides of a membrane, the solutions are **isotonic** ("same strength").

7-3 Cell Boundaries 🕩 Osmosis

Osmotic Pressure

Osmosis exerts a pressure known as osmotic pressure on the hypertonic side of a selectively permeable membrane.

7-3 Cell Boundaries 🛶 Osmosis

Because the cell is filled with salts, sugars, proteins, and other molecules, it will almost always be hypertonic to fresh water.

If so, the osmotic pressure should produce a net movement of water into the cell. As a result, the volume of the cell will increase until the cell becomes swollen or bursts.

7-3 Cell Boundaries 🕩 Osmosis

Cells in large organisms are not in danger of bursting because they are bathed in fluids, such as blood, that are isotonic.

Other cells are surrounded by tough cell walls that prevent the cells from expanding even under tremendous osmotic pressure.

7-3 Cell Boundaries 🔿 Facilitated Diffusion

Facilitated Diffusion

Cell membranes have protein channels that act as carriers, making it easy for certain molecules to cross.



7-3 Cell Boundaries IFacilitated Diffusion





7-3 Cell Boundaries 🔿 Facilitated Diffusion

Although facilitated diffusion is fast and specific, it is still diffusion.

Therefore, facilitated diffusion will only occur if there is a higher concentration of the particular molecules on one side of a cell membrane as compared to the other side.



Active Transport

Sometimes cells move materials in the opposite direction from which the materials would normally move—that is against a concentration difference. This process is known as **active transport**.

Active transport requires energy.

7-3 Cell Boundaries 📫 Active Transport

Molecular Transport

In active transport, small molecules and ions are carried across membranes by proteins in the membrane.

Energy use in these systems enables cells to concentrate substances in a particular location, even when diffusion might move them in the opposite direction.





7-3 Cell Boundaries 🛶 Active Transport

Endocytosis and Exocytosis

Large molecules and even solid clumps of material may undergo active transport by means of the cell membrane.

Endocytosis is the process of taking material into the cell by means of infoldings, or pockets, of the cell membrane.

The pocket breaks loose from the outer portion of the cell membrane and forms a vacuole within the cytoplasm.

7-3 Cell Boundaries 🔿 Active Transport

Two examples of endocytosis are:

- phagocytosis
- pinocytosis

7-3 Cell Boundaries 🔿 Active Transport

In **phagocytosis**, extensions of cytoplasm surround a particle and package it within a food vacuole. The cell then engulfs it.

Phagocytosis requires a considerable amount of energy.



7-3 Cell Boundaries 🛶 Active Transport

Exocytosis

Many cells also release large amounts of material from the cell, in a process called exocytosis.

During **exocytosis**, the membrane of the vacuole surrounding the material fuses with the cell membrane, forcing the contents out of the cell.



7-3 Section QUIZ		
_		
Continue to:		Click to Launch:
Section OUIZ	- or -	ExamView®
		Quicklake
PEARSON		Slide 40 of 47
Prentice Hall	Copyright Pearson Prentice H	al



7-3 Section QUIZ

Inlike a cell wall, a cell membrane

- a. is composed of a lipid bilayer.
- b. provides rigid support for the surrounding cell.
- c. allows most small molecules and ions to pass through easily.
- d. is found only in plants, fungi, algae, and many prokaryotes.

7-3 Section QUIZ

- 2 The concentration of a solution is defined as the
 - a. volume of solute in a given mass of solution.
 - b. mass of solute in a given volume of solution.
 - c. mass of solution in a given volume of solute.
 - d. volume of solution in a given mass of solute.

Slide

7-3 Section QUIZ

- If a substance is more highly concentrated outside the cell than inside the cell and the substance can move through the cell membrane, the substance will
 - a. move by diffusion from inside the cell to outside.
 - b. remain in high concentration outside the cell.
 - c. move by diffusion from outside to inside the cell.
 - d. cause water to enter the cell by osmosis.

7-3 Section QUIZ

4 The movement of materials in a cell against a concentration difference is called

- a. facilitated diffusion.
- b. active transport.
- c. osmosis.
- d. diffusion.

7-3 Section QUIZ

- 5 The process by which molecules diffuse across a membrane through protein channels is called
 - a. active transport.
 - b. endocytosis.
 - c. facilitated diffusion.
 - d. osmosis.